

1

JC13 Rec'd PCT/PTO 26 MAY 2009

GINSENG FERMENTED BY LACTIC ACID BACTERIUM, YOGHURT
CONTAINING THE SAME, AND LACTIC ACID BACTERIA USED IN THE
PREPARATION THEREOF

5 **TECHNICAL FIELD**

This invention relates to lactic fermenting products of ginseng obtained by fermentation of ginseng using lactic acid bacteria, yoghurt containing said lactic fermenting products of ginseng, and lactic acid bacteria used in the preparation of said lactic fermenting products of ginseng.

10 **BACKGROUND OF THE INVENTION**

Ginseng is a perennial plant belonging to the genus of Ginseng, the family of Araliaceae in the classification of the plant. About eleven kinds of ginseng have yet been known on the earth. The representative species of ginseng include Panax ginseng C. A. Meyer which grows naturally in the area of far-east Asia at 33° through 48° North Latitude (Korea, Northern Manchuria, an area of Russia) and which has excellent pharmaceutical effects; Panax quinquefolium L. which grows naturally or has been cultivated in the United States and Canada; Panax notoginseng F. H. which grows naturally or has been cultivated in China (the south-eastern area of Yunnan province and the south-western area of Guangxi province); and Panax japonicus C. A. Meyer which ranges widely in the area of Japan, the south-eastern China and Nepal.

Ginseng was classified as a top grade in plant efficacy by SINON HERBAL (Shen Nong Ben Cao Jing, called in China), the earliest herbal in China, and has been used as a valuable restorative for a long time. From many pharmacological experiments, it is demonstrated that ginseng strengthens the non-specific resistance of a human body in respect to the stress and also has an anti-acidic action. It has also been clarified that ginseng has other pharmacological effect such as improving hypertension, strengthening insulin action, lowering blood sugar level in Alloxan diabetic mouse, hepatic RNA synthesis in white rat, promoting protein synthesis and sugar and lipid metabolism together with an anti-tumor effect.

In Asian countries of Korea, China and Japan, ginseng has been used as a natural medicine for treating various diseases such as mental diseases, disorders in nervous system and diabetes.

Saponin, a principal active component of ginseng, has been known as having the effect of strengthening robustness or stamina, sedation and anti-hypertension.

At the present time, ginseng is used in form of white ginseng processed by drying crude ginseng from the cultivated at ambient temperature, red ginseng processed by heating crude ginseng (green ginseng) at 98 to 100 °C, or super red ginseng prepared by heating crude ginseng at 120 to 180 °C.

On the other hand, ginseng root contains about 4 to 10% of ginseng saponin (e.g, Panax ginseng C. A. Meyer contains 4 to 8% of ginseng saponin, and Panax quinquefolium L. contains 4 to 10% of ginseng saponin). Said ginseng saponin refers to a mixture of various ginsenosides. In particular, Panax ginseng C. A. Meyer contains relatively high content of ginsenoside Rb1, Rc and Rg1, and Panax quinquefolium L. contains relatively high content of ginsenoside Rb1 and Re.

Specific type of ginsenoside included in ginseng and their pharmacological effects are shown in the following Table 1.

Table 1

Type of Ginsenoside	Effect
Ginsenoside-Rb1	Suppression of central nerve with sedation, Suppression of central nervous voracity, Suppression of aggressive movement, Alleviation of pain, Anti-convulsion, Anti-apprehension, Accelerating secretion of adrenal cortex stimulating hormone and corticosteron, Promoting bio-synthesis of cholesterol, Improvement of memory, Lowering hyper-cholesterol, neutral fat and free aliphatic acid, Accelerating subsistence of nerves cells, Protection from liver damage, Accelerating synthesis of DNA, RNA, protein and lipid in marrow cells, Accelerating acetylcholin release, Vasodilation, Suppression of agglutination of thrombocytes, Suppression of lipid peroxidation, Promoting cholesterol metabolism, Anti-inflammation, Activation of voracity function, Suppression of hypertrophy of glomerulus.

Ginsenoside-Rb2	Accelerating sugar and fat metabolism, Anti-diabetic action, Controlling equilibrium of fat metabolism, Accelerating protein and fat synthesis, Lowering hyper-cholesterol and preventing arteriosclerosis, Antagonism on cancer toxin hormone, Suppression of proliferation of smooth muscle cells, Accelerating secretion of adrenal cortex stimulating hormone and corticosteron, DNA, RNA, Improvement of stressful decrease of appetite, Suppression of generation of tumoral vessel, Accelerating production of anti-peroxidative substance, Activation of ATP supply in liver tissues, Adjustment of immunity, Promoting cholesterol metabolism, Proliferation of liver cells and Acceleration of DNA synthesis, Suppression of agglutination of thrombocytes, and Alleviation.
Ginsenoside-Rc	Accelerating synthesis of RNA, serum cholesterol in liver, Accelerating synthesis of DNA, RNA, protein and lipid in marrow cells, Pain alleviation. Accelerating secretion of corticosteron, Promoting bio-synthesis of prostacycline and Suppression of hypertrophy of glomerulus.
Ginsenoside-Rd	Accelerating secretion of adrenal cortex stimulating hormone and corticosteron, Suppression of the hypertrophy of glomerulus.
Ginsenoside-Re	Accelerating secretion of adrenal cortex stimulating hormone and corticosteron, Alleviation of pain, Vasodilation, Anti-high temperature stress, Suppression of the proliferation of smooth muscle cells, Accelerating synthesis of DNA, RNA, protein and lipid in marrow cells, Protection from liver damage, and Promoting cholesterol metabolism.
Ginsenoside-Rg1	Strengthening immune function, Suppression of agglutination of thrombocytes, Anti-thrombin, Activation for the efficiency, Increasing an ability of memory and study, Anti-fatigue, Anti-stress, Excitation of central nerve, Vasodilation, Anti-inflammation, Anti-nephritis and functioning as a agent to increase the amount of blood stream in kidney, Function to protect from the injurious stimulation like high temperature circumstance and heat resistant-pyrogenic substance, Improvement of stressful lesion of slow motion, Accelerating subsistence of nerves cells, Proliferation of liver cells and Acceleration of DNA synthesis, Accelerating secretion of adrenal cortex stimulating hormone, Promoting cholesterol metabolism and Protection from liver damage.
Ginsenoside-Rh1	Suppression of experimental liver damage, Promotion of differentiation of tumoral cells, Suppression of agglutination of thrombocytes, Anti-inflammation and Anti-allergy.

Ginsenoside-Rh2	Suppression of cancer cell proliferation, Inducing re-differentiation of cancer cell, Suppression of permeation of cancer cells, Suppression of tumor proliferation, Increasing anti-cancer activity of an anti-cancer medicine, Anti-allergy.
Compound K	Suppressing significant generation of tumoral vessel and spread of cancer cell, Blocking secretion of IV type collagenase, Activating to form anti-regeneration vessel, and Suppressing agglutination of thrombocytes, Anti-allergy and Anti-inflammation.

It has been found that the principal active components responsible for pharmacological effects of ginseng are saponins such as ginsenoside Rb1, Rb2 and Rc. However, the active materials substantially having anti-tumor activities, capacities for suppressing metastasis of cancer cells, and/or anti-allergic effects include saponins such as Compound K (20-O- β -D-glucopyranocyl-20(S)-protopanaxidiol) as a intestinal-bacterial fermenting product, ginsenoside Rh1 and Rh2 and Δ^{20} -ginsenoside Rh2, but these saponins are contained in ginseng in extremely small amount.

Accordingly, it is necessary to increase amounts of saponins such as Compound K, ginsenoside Rh1 and Rh2 and Δ^{20} -ginsenoside Rh2 (i.e. the mixture of ginsenoside Rk2 and ginsenoside Rh3) as present at extremely lower level in the crude ginseng, in order to improve anti-cancer activities, anti-allergic effects and reinforcement of immune activity in human bodies by pharmacological effects of ginseng.

SUMMARY OF THE INVENTION

Thus, the inventors have also made attempts to more efficiently obtain saponins such as Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2. As a result, the inventors has found that lactic fermenting products of ginseng obtained by fermentation of ginseng using lactic acid bacteria contain significantly more amounts of Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2. The present invention is based on such findings.

Accordingly, it is one object of the invention to provide lactic fermenting products of ginseng.

Further, it is other object of the invention to provide yoghurts containing said lactic fermenting products of ginseng.

Further, it is another object of the invention to provide lactic acid bacteria used for the preparation of said lactic fermenting products of ginseng.

DETAILED DESCRIPTION OF THE INVENTION

5 According to the invention, (1) lactic fermenting products of ginseng, (2) yoghurts containing said lactic fermenting products of ginseng, and (3) lactic acid bacteria advantageously used for the preparation of said lactic fermenting products of ginseng, are provided.

The invention is described more specifically here in below.

10 According to the invention, when ginseng is fermented by lactic acid bacteria, the saponin ingredients of ginseng prior to fermentation are to be bio-converted to the saponin ingredients of Compound K(20-O- β -D-glucopyranocyl-20(S)-protopanaxadiol), ginsenoside Rh1 and ginsenoside Rh2, and Δ^{20} -ginsenoside Rh2 which were not included at all or included infinitesimally in said ginseng prior to the fermentation. Then, lactic fermenting products of
15 ginseng comprise at least one ingredient selected from the group consisting of Compound K(20-O- β -D-glucopyranocyl-20(S)-protopanaxadiol), ginsenoside Rh1 and ginsenoside Rh2, and Δ^{20} -ginsenoside Rh2.

According to general knowledge in the relevant art, the saponin ingredients of Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2 have more excellent activity of anti-
20 cancer and anti-allergy or suppressing cancer spread as compared with the same of ginsenoside Rb1, Rb2 and Rc(see the literatures of "Bae et. al., Biol. Pharm. Bull., 25, 743-747, 2002 ; Bae et al., 25, 58-63, 2002 ; Wakabayashi et al., Oncol. Res., 9, 411-417, 1998 ; Saiki et. al., Proceedings of the 8th international symposium on Ginseng(Korea Ginseng Academy, Seoul, Korea), 305-316, 2002 ; Hasegawa and Saiki, Proceedings of the 8th international symposium
25 on Ginseng(Korea Ginseng Academy, Seoul, Korea), 317-334, 2002).

On the other hand, sources of ginseng which can be used for the fermentation by lactic acid bacterium are not particularly limited and the Korean ginseng (*Panax ginseng* C. A. Meyer) as well as other kinds of ginsengs such as *Panax quinquefolium* L., *Panax Notoginseng* F. H. Chen and *Panax japonicus* C. A. Meyer in their natural form or their processed product
30 can be employed. More specifically, either at least one of the ginseng leaf, ginseng extract

and ginseng powder, or green ginseng, red ginseng, white ginseng and other kinds of ginsengs afore-mentioned(hereinafter inclusively referred to as "ginseng") can be employed. In terms of processing, the treatment with acid, at high temperature and under pressure can be desirably applied to give lactic fermenting products of ginseng.

5 Pulverizing degree of ginseng in a form of dried powder is not particularly restricted. Ginseng can be pulverized so that lactic acid bacterium may penetrate very effectively into tissues or fibrous cells of ginseng. Such pulverizing degree of ginseng and other pulverization method has been commonly known to a person in the pertinent art. Though the treatment with acid, at high temperature and pressure can be readily applied to a product *per se*, it is preferable
10 to adopt the treatment of dry pulverization in consideration of the effect of the treatment and the fermentation efficiency thereafter.

Pulverizing degree of ginseng in the dry pulverization is not particularly restricted too.

In the case that at least one pulverized ginseng material selected from the group consisting of the green ginseng, red ginseng, white ginseng, fine ginseng, *Panax quinquefolium* L.,
15 ginseng leaf, ginseng extract and ginseng powder (hereinafter, "the ginseng source") is fermented by lactic acid bacterium, the amount of Compound K is particularly increased in the product of ginseng fermented by lactic acid bacterium.

The ginseng product by acid treatment can be obtained by adding an acid, preferably acetic acid, lactic acid, citric acid, butyric acid, tartaric acid, propionic acid or hydrochloric acid
20 to at least one pulverized ginseng material selected from the group consisting of the green ginseng, red ginseng, white ginseng, fine ginseng, *Panax quinquefolium* L., ginseng leaf, ginseng extract and ginseng powder and cultivating the mixture at 60°C for 5 hours and then neutralizing with calcium salt. According to the invention, in the case that ginseng treated with an acid is fermented by lactic acid bacterium, the amounts of ginsenoside Rh1 and Rh2 are
25 particularly increased in the resultant fermenting products by lactic acid bacterium.

The ginseng product can be prepared by treating ginseng at high temperature, e.g., red ginseng can be prepared by heating said pulverized ginseng material at 100°C for 2 to 5 hours. According to the invention, in case that ginseng treated at high temperature is fermented by lactic acid bacterium, the amounts of Compound K and ginsenoside Rh1 and Rh2 are
30 particularly increased in the resultant fermenting products by lactic acid bacterium.

The ginseng treated under pressure can be prepared by treating said pulverized ginseng material under pressure at 110 to 130°C for 2 to 5 hours. According to the invention, in case that ginseng treated under pressure is fermented by lactic acid bacterium, the amounts of ginsenoside Rh1 and Rh2 and Δ^{20} -ginsenoside Rh2 are particularly increased in the resultant lactic fermenting products of ginseng despite the low amount of Compound K.

On the other hand, the method for preparing lactic fermenting products of ginseng is not particularly limited and can be carried out by a method employed generally in the pertinent art under the proper condition for the fermentation with lactic acid bacterium.

Specifically, lactic fermenting products of ginseng can be prepared through the suspension of the pulverized ginseng material in water for lactic acid bacterium to be cultured at suitable temperature for 48 to 72 hours and passing through a process of bio-converting for the resultant lactic fermenting products of ginseng, centrifuging the resultant product and then filtrating the upper transparent liquid only and concentrating the liquid to give lactic fermenting products of ginseng.

The lactic acid bacterium used for the fermentation is not particularly limited, as long as the bio-converting rate to Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2 is high. For instance, bacteria of *Lactobacillus*, *Streptococcus* or *Bifidobacterium* spp. can be used. More specifically, at least one of *Bifidobacterium* K-103(see Arch. Pharm. Res., 21, 54-61, 1998, Professor Dong-Hyun, Kim, the college of pharmacy, Kyung-Hee University, Seoul, Korea), *Bifidobacterium* K-506(see Arch. Pharm. Res., 21, 54-61, 1998, Professor Dong-Hyun, Kim, the college of pharmacy, Kyung-Hee University, Seoul, Korea), *Bifidobacterium* cholerium KK-1(KCCM-10364), *Bifidobacterium* minimum KK-2(KCCM-10365), *Bifidobacterium* H-1(KCCM-10493) and *Bifidobacterium* KK-11(Professor Dong-Hyun, Kim, the college of pharmacy, Kyung-Hee University) can be employed.

Among these, *Bifidobacterium* H-1, *Bifidobacterium* cholerium KK-1 and *Bifidobacterium* minimum KK-2 are the microorganisms which have been firstly developed by the inventors to obtain lactic fermenting products of ginseng. Said bifadobacterium cholerium KK-1, *Bifidobacterium* minimum KK-2 and *Bifidobacterium* H-1 have been deposited respectively as the Accession numbers of KCCM-10364(March 22, 2003), KCCM-10365(March 22, 2003), KCCM-10493(May 1st, 2003) with the Korean Culture Center of

Microorganism. *Bifidobacterium cholerae* KK-1 and *Bifidobacterium minimum* KK-2, *Bifidobacterium* H-1 and *Bifidobacterium* K-11 of the invention are gram positive and anaerobic, and also fructose 6-phosphate phosphoketolase positive.

The bacilli have the sugar usability as shown in the following table.

5

CLASS	Sugar Usability			
	<i>Bifidobacterium</i> KK-1	<i>Bifidobacterium</i> KK-2	<i>Bifidobacterium</i> H-1	<i>Bifidobacterium</i> KK-11
amigladin	-	-	-	-
arabinose	-	+/-	+	+
cellobiose	-	+/-	-	-
dextrin	+	+	-	-
esculin	-	-	+/-	+/-
fructose	-	+	+	+/-
galactose	+	+	+	+
gluconate	-	-	-	-
glucose	+/-	+	+	+
glycogen	+/-	+	-	-
inositol	-	-	-	-
inulin	-	+/-	+/-	+/-
lactose	+	+/-	+	+
maltose	+/-	+	+	+
mannitol	-	+	+	+
mannose	+/-	+	+	+
melezitose	-	-	-	-
melibiose	+	-	+	+
raffinose	+	-	+	+
ribose	+/-	+/-	+/-	+/-
salicin	+	+	+	+
sorbitol	+/-	-	-	-
starch	+	+	-	-
sucrose	-	+/-	+	+
trehalose	-	-	-	-
xylose	-	+/-	+	+

Said *Bifidobacteria* KK-1, KK-2, H-1 and K-11 are characteristically identical with the

other bacterium of the same species in the classification. However, the Bifidobacteria enables to provide lactic fermenting products of ginseng thereby which includes much higher amount of Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2, when applied to the processes for fermentation of ginseng.

5 Accordingly, the lactic fermenting products of ginseng of the prevent invention contains much higher amount of Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2. Particularly, it is preferable that the whole amount of (Compound K+ginsenoside Rh1), (ginsenoside Rh1+ginsenoside Rh2), (ginsenoside Rh2+ Δ^{20} -ginsenoside Rh2+ginsenoside Rh1) or (Compound K+ginsenoside Rh1+ginsenoside Rh2) be included respectively in the ratio
10 of more than 0.1 with respect to total amount of (ginsenoside Rc+ginsenoside Rd+ginsenoside Rb1+ginsenoside Rb2+ginsenoside Re+ginsenoside Rg1). It is further preferable that the ratio of the both components be in the range of 1 : 50 to 50 : 1 in case of the production of (Compound K+ginsenoside Rh1) and 50 to 50 : 1 in case of the production of (ginsenoside Rh2+ginsenoside Rh1) and that the ratio of the sum of (ginsenoside Rh2+ Δ^{20} -ginsenoside Rh2) : ginsenoside Rh1 be in the range of 1 : 50 to 50 : 1 in case of the production of (ginsenoside Rh2+ Δ^{20} -ginsenoside Rh2+ginsenoside Rh1) and the ratio of Compound K : ginsenosides be in the range of 1 : 50 to 50 : 1 in case of the production (Compound K+ginsenoside Rh1+ginsenoside Rh2). Not included naturally in red ginseng, Compound K appears increasingly in red ginseng, when fermented by lactic acid bacterium according to the
15
20 invention.

Said lactic acid bacteria of the invention appear to suppress the intestine-harmful-bacillus and/or to suppress the activity of the intestine-harmful-enzyme and/or to suppress the proliferation of cancer cells.

On the other hand, the invention provides also a ginseng yoghurt containing the resultant
25 substance of the fermentation by lactic acid bacterium.

As said ginseng yoghurt contains the resultant lactic fermenting products of ginseng according to the invention, namely, includes plenty of the ginseng saponin ingredients of Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2, the yoghurt of the invention is certainly a multi-functional yoghurt to exert the high physiological function of
30 Compound K, ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2, more specifically the

functions of anti-cancer, anti-allergy and the reinforcement of immunity. Such ginseng yoghurt can be prepared by merely adding 0.1 to 10% by weight of said resultant lactic fermenting products of ginseng to a usual yoghurt. However, this method requires separately a process for the fermentation of milk of yoghurt and another process for the additive
5 fermentation of ginseng by lactic acid bacterium to obtain lactic fermenting products of ginseng. Thus, this method is not industrially favorable for economic reasons. Accordingly, the ginseng yoghurt of the invention can be prepared by fermenting concurrently milk of yoghurt and 0.1 to 10% by weight of ginseng by lactic acid bacterium. In the reactant for the fermentation, the vitamins can be included.

10 The bacterium to prepare a ginseng yoghurt is not particularly limited, if enabling to ferment concurrently both ginseng and milk for the preparation of yoghurt. For instance, the bacterium of Lactobacillus, Streptococcus bacillus or Bifidobacterium, preferably at least one selected from the group of Bifidobacterium K-103, Bifidobacterium K-506, Bifidobacterium cholerium KK-1, Bifidobacterium minimum KK-2, Bifidobacterium H-1 and Bifidobacterium
15 KK-II can be used.

Milk for preparing yoghurt is not particularly restricted, and goat milk, sheep milk, nonfat milk, skim milk as well as cow milk can be used.

After the fermentation to prepare a yoghurt, lactic acid bacterium is optionally removed. However, as lactic acid bacterium has a favorable function to improve the intestinal conditions,
20 it is preferable not to remove said lactic acid bacterium. Accordingly, the yoghurt of the invention can improve the intestinal conditions.

On the other hand, the ginseng powder yoghurt can be available by adding ginseng powder to the resultant yoghurt. Accordingly, the yoghurt is not particularly limited to a formal class. Then, various types of usual yoghurt prepared through the fermentation of milk by lactic acid
25 bacterium can be possible.

The invention will be understood more readily with reference to the following examples, however, these examples are intended to illustrate the invention only and are not to be construed to limit the scope of the invention. The modification and application thereof usually acceptable in the pertinent art fall within the scope of the invention.

THE PREFERRED EMBODIMENTS OF THE INVENTION**Example A1**

Undried ginseng root (5-year *P. ginseng* C. A. Meyer root from Kumsan in Korea procured from Kyungdong Market) was washed thoroughly with hot water, dried and ground to a fine powder. Subsequently 1g of powdered dry ginseng prepared as described above and 0.1g of vitamin C were suspended in 100ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24hrs at 37°C.

10 Example A2

Each 2g of powdered dry ginseng prepared from sufficiently dehydrated undried ginseng root or the root hair of ginseng and 0.1g of vitamin C were suspended in 100ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* K-103 and *Bifidobacterium* K-506 thereto, followed by a fermentation for 24hrs at 37°C.

15

Example A3

1g of powdered dry ginseng prepared from sufficiently dehydrated undried ginseng root and 0.1g of vitamin C were suspended in 100 ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Lactobacillus bulgaricus*, *Streptococcus thermophilus*, *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 12hrs at 37°C.

20

Example A4

1g of powdered dry white ginseng root and 0.1g of vitamin C were suspended in 100 ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Bifidobacterium* KK-1 thereto, followed by a fermentation for 24 hrs at 37°C.

25

Example A5

1g of powdered dry ginseng root and 0.1g of vitamin C were suspended in 100 ml of milk.

30

Ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24hrs at 37°C.

5 Example A6

Powdered dry ginseng root was treated with 0.5% lactic acid at 60°C for 5hrs followed by a neutralization and drying. 1g of powdered dry ginseng root (acid-treated ginseng) prepared as described above and 0.1g of vitamin C, were suspended in 100ml of milk. Acid-treated ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated
10 *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24hrs at 37°C.

Example A7

Powdered dry ginseng root was suspended in distilled water and steamed at 100°C for 2hrs
15 followed by a drying. 1g of powdered dry ginseng root (ginseng treated at high temperature) prepared as described above and 0.1g of vitamin C were suspended in 100ml of milk. high temperature treated ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24 hrs at 37°C.

20

Example A8

Powdered dry ginseng root was pressurized for 2hrs at 120°C followed by a drying. 1g of powdered dry ginseng root (pressurized ginseng) prepared as described above and 0.1g of vitamin C were suspended in 100ml of milk. Pressurized ginseng yogurt was then obtained by
25 inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24hrs at 37°C.

Example A9

1g of powdered dry ginseng root and 0.1g of vitamin C were suspended in 100 ml of milk.
30 Ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated

Bifidobacterium KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24hrs at 37°C.

Example A10

5 Powdered dry ginseng root was treated with 5% lactic acid at 60°C for 5hrs followed by a neutralization and drying. 1g of powdered dry ginseng root (high temperature treated ginseng) prepared as described above and 0.1g of vitamin C were suspended in 100ml of milk. high temperature treated ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed
10 by a fermentation for 24 hrs at 37°C.

Example A11

Powdered dry ginseng root was suspended in distilled water and steamed at 100°C for 2hrs followed by a drying. 1g of powdered dry ginseng root (ginseng treated at high temperature)
15 prepared as described above and 0.1g of vitamin C were suspended in 100ml of milk. high temperature treated ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24 hrs at 37°C.

20 Example A12

Powdered dry ginseng root was pressurized at 120°C for 2 hrs followed by a drying. 1g of powdered dry ginseng root (pressurized ginseng) prepared as described above and 0.1g of vitamin C were suspended in 100ml of milk. Pressurized ginseng yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and
25 *Bifidobacterium* KK-2 thereto, followed by a fermentation for 24hrs at 37°C.

Examples A13-A24

Each of lactic acid bacteria-fermented products was obtained by repeating exactly the same
30 process as examples A1 to A12 above, except 2g (wet weight) of precultivated lactic acid

bacteria and 100 ml of distilled water instead of 100ml of milk were used.

Experimental example A1: amount analysis of saponin component

1g of commercially available regular ginseng root (Nonghyup 4-6 year *P. ginseng* C. A. Meyer
5 root from Kumsan in Korea procured from Kyungdong Market) and each 1 g of its powdered
dry, acid-treated, high temperature treated and pressurized version(prepared by the same
process as in examples above) was individually mixed with 100ml of milk containing 0.1g of
Vitamin C. Each of the above mixture was subsequently inoculated with 1 g of
Bifidobacterium KK-1 and *Bifidobacterium* KK-2, and fermented for 72hrs at 37°C, followed
10 by a concentration by decompression to obtain fermented yogurt of ginseng, high temperature
treated ginseng, acid-treated ginseng and pressurized ginseng, respectively. Subsequently, 2 g
of ginseng and each 2 g of its powdered dry, acid-treated, high temperature treated and
pressurized version, and each 2 g of fermented products above were extracted with 100ml of
methanol 3 times. Each of the extracts was then concentrated and suspended in water followed
15 by a extraction with 100ml of ether 3 times. Each of the ether fractions was further extracted
with 100ml of butanol 3 times. The butanol fractions were then subject to concentration and
subsequent TLC analysis after dissolved in methanol (Solvent system:
 $\text{CHCl}_3:\text{MeOH}:\text{H}_2\text{O}=65:35:10/\text{CHCl}_3:\text{EtOAc}:\text{MeOH}:\text{H}_2\text{O}=15:40:22:9$; Spraying reagent: 5%
sulfuric acid in methanol; TLC scanner: Shimadzu CS-9301PC). The results obtained are
20 shown in Table 2. The amount of each component was calculated based on which is contained
in 100% of the finally extracted fraction.

Similar results as in Table 2 were also obtained with examples A1 to A4 and A9 to A12.
Fermented ginseng yogurts according to Examples A13 to A24 were shown to contain similar
25 amount of saponin to that of fermented yogurts as described above.

Table 2

Name of the components	amount of component (%)			
	Ginseng (example A5)	fermented yogurt of ginseng (example A5)	high temperature treated ginseng (example A7)	fermented yogurt of high temperature treated ginseng (example A7)
ginsenoside Rb1	15.1	1.6	5.1	2.7
ginsenoside Rb2	8.2	1.1	3.5	2.1
ginsenoside Rc	9.5	0.5	3.8	2.9
ginsenoside Re	10.7	2.7	7.8	4.5
ginsenoside Rg3	< 1	< 1	14.6	8.5
20-ginsenoside Rg3	< 1	< 1	4.5	< 1
compound K	0	28.6	< 1	2.9
ginsenoside Rh2	< 1	< 1	< 1	3.2
Δ^{20} -ginsenoside Rh2	< 1	< 1	< 1	< 1
ginsenoside Rh1	< 1	1.2	0.5	1.8
Protopanaxadiol	< 1	2.1	< 1	2.1
Name of the components	amount of component (%)			
	acid-treated ginseng (example A6)	fermented yogurt of acid- treated ginseng (example A6)	pressurized ginseng (example A8)	fermented yogurt of pressurized ginseng (example A8)
ginsenoside Rb1	2.5	1.2	2.5	1.2
ginsenoside Rb2	2	0.9	1.2	0.8
ginsenoside Rc	1.8	1.1	1.5	1.0
ginsenoside Re	6.8	5.2	3.9	2.9
ginsenoside Rg3	25	7	16.1	4.5
20-ginsenoside Rg3	< 1	< 1	6.5	2.1
compound K	0	2.2	0	1.2
ginsenoside Rh2	0.2	10.2	< 1	4.5
Δ^{20} -ginsenoside Rh2	0.1	1.8	< 1	3.8
ginsenoside Rh1	0.5	2.1	2.1	2.5
protopanaxadiol	< 1	2.5	< 1	2.8

Experimental example A2: analysis of anticancer effect

HepG2 (Human liver cancer cell line; KCLB-10023), A-549 (Human lung cancer cell line; KCLB-10185), P-388(mouse lymph neoplastic cell line; KCLB-10046) and L-1210 (mouse lymphocytic leukemia cell line; KCLB-10219) were each grown in RPMI 1640 medium supplemented with 10% FBS, 1% antibiotics-antimycotics(GIBCO,USA) and 2.2g of NaHCO₃. Subsequently, HepG2 and A-549 cells were harvested by trypsinization (0.25% of trypsin) and each 180 μ l of cells thereof was seeded into each well of a 96 well plate at 3 x 10⁴ cells/well. Then the plates were incubated for 24hrs at 37 $^{\circ}$ C in 5% CO₂ incubator. For P-388 and L-1210 cells, each 180 μ l of cells harvested by trypsinization (0.25% of trypsin) was seeded into each well of a 96 well plate at 4 x 10⁴ cells/well. Then the plates were incubated for 2 hrs at 37 $^{\circ}$ C in 5% CO₂ incubator. Butanol extracts of white ginseng and lactic acid bacteria fermented ginseng as in Table 3 were autoclaved and 20 μ l of each was added to each well of 96 well plate prepared as described above at the concentration of 10mg/ml. The 96 well plates were then incubated for 48hrs at 37 $^{\circ}$ C in 5% CO₂ incubator. After 48 hrs, 20mg/ml of MTT was added to each well of the 96well plates and they were further incubated for 4hrs in 5% CO₂ incubator. After 4hrs, the media were removed from each well of the 96 well plates and 100 μ l of DMSO was added thereto. Subsequently, the absorbance at 580nm was determined for each well using ELISA reader to test for cell toxicity. The results obtained are shown in Table3.

Table 3

Type of ginseng	ED50(μ g/ml)			
	P388	L1210	A549	HepG2
extract of regular ginseng	> 100	> 100	> 100	> 100
lactic acid bacteria - fermented product of white ginseng	98	50	160	96
lactic acid bacteria- fermented product of acid- treated ginseng	82	51	102	95
lactic acid bacteria- fermented product of high temperature treated ginseng	78	45	87	91

lactic acid bacteria- fermented product of pressurized ginseng	85	52	105	92
----------------------------------------------------------------------	----	----	-----	----

Experimental example A3 : The effect on *E.Coli* HGU-3 and harmful intestinal enzymes

Fermented ginseng products obtained from the examples above and 24hr-cultivated *E.Coli* HGU-3 (from Dong-Hyun Kim of School of Pharmacy, Kyunghee University, and properties are almost the same as those of *E. coli* in general.) together with each of *Bifidobacterium* KK-1 or *Bifidobacterium* KK-2 at the concentration of 10^7 cells were inoculated into 5ml of GAM solution and was incubated for 24hrs. Subsequently the enzyme activities of β -glucuronidase and tryptophanase produced by bacterial strains, which are known as the cause for colon cancer and liver disorder, respectively, were measured.

The enzyme activity of β -glucuronidase was measured by adding 0.02ml of 10mM p-nitrophenyl- β -D-glucuronide and 0.1ml of enzyme solution to 0.38ml of 0.1M phosphate buffer (pH7.0) followed by an 1 hour incubation at 37°C. The reaction was then terminated by addition of 0.5ml of 0.5N NaOH, and 1ml of distilled water was added thereto followed by a centrifugation (2000 x g, 20 min). Absorbance at 405nm was then determined for each of the resulting supernatants.

The enzyme activity of tryptophanase was measured by adding 0.2ml of complete reaction mixture (0.1M bicine, pH8.0, 4% pyridoxal-5-phosphate, 20% bovine serum albumin), 0.2ml of 0.02M tryptophan and 0.1ml of enzyme solution followed by a 30min incubation. The reaction was then terminated by addition of 2ml dye solution(p-dimethylaminobenzaldehyde 14.7g, 95% ethanol 948ml, C-H₂SO₄ 52ml), and subsequently centrifuged at 2000 x g for 20min. Absorbance at 550nm was then determined for each of the resulting supernatants.

The results obtained are shown in Table 4 and Table 5 for *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2, respectively.

Table 4

	% inhibition	
	β -glucuronidase	tryptophanase
control (KK-1 and total intestinal flora cultivated in GAM)	0	0
ginseng extract (KK-1 and total intestinal flora cultivated in GAM containing 1% of ginseng extract)	32	45
lactic acid bacteria-fermented product of ginseng (KK-1 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria-fermented product of ginseng)	85	85
lactic acid bacteria-fermented product of acid treated ginseng (KK-1 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of acid-treated ginseng)	75	78
lactic acid bacteria-fermented product of high temperature treated ginseng (KK-1 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of high temperature treated ginseng)	89	87
lactic acid bacteria-fermented product of pressurized ginseng (KK-1 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of pressurized ginseng)	78	85

Table 5

	% inhibition	
	β -glucuronidase	tryptophanase
control (KK-2 and total intestinal flora cultivated in GAM)	0	0
Ginseng extract (KK-2 and total intestinal flora cultivated in GAM containing 1% of ginseng extract)	32	45
lactic acid bacteria fermented product of ginseng (KK-2 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of ginseng)	76	81
lactic acid bacteria fermented product of acid treated ginseng (KK-2 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of acid-treated ginseng)	67	88
lactic acid bacteria fermented product of high temperature treated ginseng (KK-2 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of high temperature treated ginseng)	82	68
lactic acid bacteria fermented product of pressurized ginseng (KK-2 and total intestinal flora cultivated in GAM containing 1% of lactic acid bacteria fermented product of pressurized ginseng)	71	78

It is known from Table 4 and 5 that the fermented ginseng products when absorbed into cells, have anticancer effect, suppress the activity of β -glucuronidase and tryptophanase produced by intestinal bacteria, and thus have preventing effect on colon cancer and liver damage.

Example B1

1g of *Panax quinquefolium* L. root as powdered (can contain 0.1g of vitamin C) was suspended in 100 ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of

precultivated lactic acid bacteria *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Bifidobacterium* KK-1 thereto, followed by a incubation for 24hrs at 37°C.

Example B2

- 5 1g of *Panax quinquefolium* L. root as powdered (can contain 0.1g of vitamin C) was suspended in 100 ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Lactobacillus bulgaricus*, *Streptococcus thermophilus* and *Bifidobacterium* KK-1 thereto, followed by a incubation for 24hrs at 37°C.

10 Example B3

1g of *Panax quinquefolium* L. root as dry-powdered (can contain 0.1g of vitamin C) was suspended in 100 ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-2 thereto, followed by a incubation for 24hrs at 37°C.

15

Example B4

- Panax quinquefolium L. root as dry-powdered was suspended in water and lactic acid was added thereto at the final concentration of 1%. Then it was incubated at 60°C for 5hrs followed by a neutralization and drying. 1g of powdered dry ginseng root (can contain 0.1g of vitamin C) prepared as described above was suspended in 100ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-11 thereto, followed by a incubation for 24hrs at 37°C.
- 20

Example B5

- 25 1 g of *Panax quinquefolium* L. root as dry-powdered was moisturized and steamed for 2hrs (can contain 0.1g of vitamin C). And it was suspended in 100 ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-11 thereto, followed by a incubation for 24hrs at 37°C.

30 Example B6

Panax quinquefolium L. root as dry-powdered (can contain 0.1g of vitamin C) was pressurized for 2hrs at 120°C and 1 g thereof was suspended in 100ml of milk. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated *Bifidobacterium* KK-1 and *Bifidobacterium* KK-11 thereto, followed by a incubation for 24hrs at 37°C.

5

Example B7

Panax quinquefolium L. root as dry-powdered (can contain 0.1g of vitamin C) was pressurized for 2hrs at 120°C and it was suspended in 100ml of milk to give final concentration of 5%. Yogurt was then obtained by inoculating each 1ml (about 10^9 cells/ml) of precultivated
10 *Bifidobacterium* KK-1 and *Bifidobacterium* KK-11 thereto, followed by a incubation for 24hrs at 37°C.

Examples B8-B14

Each of lactic fermenting products of ginseng was obtained by repeating exactly the same
15 process as examples B1 to B7 above, except 2g (wet weight) of precultivated lactic acid bacteria and 100 ml of distilled water instead of 100ml of milk were used.

Experiment example B1: Content analysis of saponin component

1g of commercially available general Panax quinquefolium L. (4-year-old Panax quinquefolium
20 L. root from Canada procured from Hongrim Trading Co., Ltd.) and each 1g of its powdered dry, acid-treated, high temperature treated and pressurized version were individually mixed with 100ml of milk containing 0.1g of Vitamin C. Each of the above mixture was subsequently inoculated with 1 g of *Bifidobacterium* KK-1 and *Bifidobacterium* KK-11 and incubated for 72hrs at 37°C followed by a concentration by decompression to obtain fermented yogurt of dry
25 Panax quinquefolium L. , high temperature treated Panax quinquefolium L. , acid-treated Panax quinquefolium L. and pressurized Panax quinquefolium L. , respectively. Subsequently, Panax quinquefolium L. and its powdered dry, acid-treated, high temperature treated and pressurized version, and each of the fermented products above (2-20g) were extracted with 100ml of methanol 3 times. Each of the extracts was then concentrated and suspended in water
30 followed by a extraction with 100ml of ether 3 times. Each of the ether fractions was further

extracted with 100ml of butanol 3 times. The butanol fractions were then subject to concentration, and subsequent TLC analysis after dissolved in methanol (Solvent system: $\text{CHCl}_3:\text{MeOH}:\text{H}_2\text{O}=65:35:10$ / $\text{CHCl}_3:\text{EtOAc}:\text{MeOH}:\text{H}_2\text{O}=15:40:22:9$; Spraying reagent: 5% sulfuric acid in methanol; TLC scanner: Shimadzu CS-9301PC). The results are shown in Table

5 6.

The amount of saponin shown in the following table was calculated based on which is in 100% of the finally extracted fraction. The control contains exactly the same products as the experimental samples except containing no ginseng.

10

Table 6

Name of the components	amount of chemical constituent(%)			
	Panax quinquefolium L.	fermented yogurt of Panax quinquefolium L. (example B3)	high temperature treated Panax quinquefolium L.	fermented yogurt of high temperature treated Panax quinquefolium L. (example B5)
ginsenoside Rb1	45.9	11.2	20.3	5.9
ginsenoside Rb2	1.1	< 1	< 1	< 1
ginsenoside Rc	4.8	< 1	1.5	< 1
ginsenoside Re	23.6	12.3	9.9	3.5
ginsenoside Rg3	< 1	< 1	19.1	4.6
20-ginsenoside Rg3	< 1	< 1	2.5	< 1
compound K	0	25.4	< 1	5.3
ginsenoside Rh2	< 1	< 1	< 1	4.6
Δ^{20} -ginsenoside Rh2	< 1	< 1	< 1	1.2
ginsenoside Rh1	< 1	2.8	4.8	1.8
protopanaxadiol	< 1	< 1	< 1	< 1

Name of the components	amount of chemical constituent(%)			
	acid-treated Panax quinquefolium L.	fermented yogurt of acid- treated Panax quinquefolium L. (example B4)	pressurized Panax quinquefolium L.	fermented yogurt of pressurized Panax quinquefolium L. (example B6)
ginsenoside Rb1	4.5	1.4	11.8	3.2
ginsenoside Rb2	< 1	< 1	1	< 1
ginsenoside Rc	< 1	< 1	< 1	< 1
ginsenoside Re	6.8	4.2	9.3	3.1
ginsenoside Rg3	28	11.5	23.2	9.6
20-ginsenoside Rg3	2.5	< 1.1	6.5	3.1
compound K	0	2.8	0	3.7
ginsenoside Rh2	0.2	11.3	< 1	8.6
Δ^{20} -ginsenoside Rh2	0.1	1.4	< 1	1.1
ginsenoside Rh1	1.5	3.7	1.2	2.5
protopanaxadiol	< 1	< 1	< 1	1.4

Experimental example B2 : Analysis of anticancer effect

HepG2 (Human liver cancer cell line; KCLB-10023), A-549 (Human lung cancer cell line; KCLB-10185), P-388(mouse lymph neoplastic cell line; KCLB-10046), L-1210(mouse lymphocytic leukemia cell line; KCLB-10219) were each grown in RPMI 1640 medium supplemented with 10% FBS, 1% antibiotics-antimycotics (GIBCO, USA) and 2.2g of NaHCO₃. Subsequently, HepG2 and A-549 cells were harvested by trypsinization(0.25% of trypsin) and each 180 μ l of cells thereof was seeded into each well of a 96 well plate at 3 x 10⁴ cells/well. Then the plates were incubated for 24hrs at 37°C in 5% CO₂ incubator. For P-388 and L-1210 cells, each 180 μ l of cells harvested by trypsinization(0.25% of trypsin) was seeded into each well of a 96 well plate at 4 x 10⁴ cells/well. Then the plates were incubated for 2hrs at 37°C in 5% CO₂ incubator. Panax quinquefolium L. extracts and lactic fermenting products of ginseng prepared as in Table 3 (Butanol extracted fraction after fermented with lactic acid bacteria) were autoclaved and 20 μ l of each was added to each well of 96 well plate as described above at the concentration of 10mg/ml. The 96 well plates were then incubated for

48hrs at 37°C in 5% CO₂ incubator. After 48 hrs, 20mg/ml of MTT was added to each well and further incubated for 4hrs in 5% CO₂ incubator. After 4hrs, the media were removed from each well of the 96 well plates and 100 μ l of DMSO was added thereto. Subsequently, the absorbance at 580nm was determined for each well using ELISA reader to test for cell toxicity.

5 The results obtained are shown in Table 7.

Table 7

Type of ginseng	ED50(μ g/ml)			
	P388	L1210	A549	HepG2
Extract of regular <i>Panax quinquefolium</i> L.	> 100	> 100	> 100	> 100
lactic fermenting products of <i>Panax quinquefolium</i> L.	78	65	> 100	89
lactic fermenting products of acid-treated <i>Panax quinquefolium</i> L.	89	65	95	85
lactic fermenting products of high temperature treated <i>Panax quinquefolium</i> L.	92	50	89	95
lactic fermenting products of pressurized <i>Panax quinquefolium</i> L.	92	60	110	95

Experimental example B3 : The effect on *E. coli* HGU-3 and harmful intestinal enzymes

10 24hr-cultivated *E. Coli* HGU-3(from Dong-Hyun Kim of School of Pharmacy, Kyunghee University and properties are almost the same as those of *E. coli* in general.) (or freshly prepared total intestinal flora of human) together with each of *Bifidobacterium* KK-1(or *Bifidobacterium* KK-2 or *Bifidobacterium* H-1 or *Bifidobacterium* KK-11) at the concentration of 10⁵ cells were inoculated into 5ml of GAM solution (the one before this contains ginseng; no
15 ginseng) and it was incubated for 24hrs. The enzyme activity of β -glucuronidase and tryptophanase produced by bacterial strains, which are known as the cause for colon cancer and liver disorder, respectively, were measured.

The enzyme activity of β -glucuronidase was measured by adding 0.02ml of 10mM p-

nitrophenyl- β -D-glucuronid and 0.1ml of enzyme solution to 0.38ml of 0.1M phosphate buffer(pH7.0) followed by 1 hour incubation at 37°C. The reaction was then terminated by addition of 0.5ml of 0.5N NaOH and 1ml of distilled water was added thereto followed by a centrifugation (2000 x g, 20 min). Absorbance at 405nm was then determined for each of the

5 resulting supernatants.

The enzyme activity of tryptophanase was measured by adding 0.2ml of complete reaction mixture (0.1M bicine, pH8.0, 4% pyridoxal-5-phosphate, 20% bovine serum albumin), 0.2ml of 0.02M tryptophan, 0.1ml of enzyme solution followed by a 30 min incubation. The reaction was then terminated by addition of 2ml dye solution(p-dimethylaminobenzaldehyde 14.7g, 95% ethanol 948ml, C-H₂SO₄ 52ml), and subsequently centrifuged at 2000 x g for 20min. Absorbance at 550nm was then determined for each of the resulting supernatants.

10

The quantitative analysis of ammonia produced was done by adding 0.1M phosphate buffer (pH7.0), 20 μ l of fractional suspension and 100 μ l of 1N H₂SO₄, and adding each 1 ml of solution mixture 1 (1% propanol, 0.005% sodium nitroprusside) and solution mixture 2 (0.1% sodium hypochlorite, 0.5% sodium hydroxide, 5.5% sodium phosphate dibasic) thereto followed by a heating at 60°C for 20min and subsequent cooling to room temperature. The absorbance at 660nm was then determined.

15

The results obtained are shown in Table 8 where the inhibitory effects on the activity of colon cancer causing enzymes are shown for the mixed culture of *E. coli* HUG-3 and lactic acid bacteria, and that of total intestinal flora of human and lactic acid bacteria(control contains only *E. coli* HUG-3).

20

Table 8

	% inhibition		
	β -glucurodinase	tryptophanase	production of ammonia
control	0	0	0
<i>Bifidobacterium</i> KK-1	85	78	67
<i>Bifidobacterium</i> KK-2	70	89	75
<i>Bifidobacterium</i> H-1	32	45	59
<i>Bifidobacterium</i> KK-11	68	75	69

It is known from Table 8 that the fermented products of ginseng, when absorbed into cells, have anticancer effect, and suppress the activity of β -glucuronidase and tryptophanase produced by intestinal bacteria and also the production of ammonia, thus have preventing effect on colon cancer and liver damage.

5

Experimental example B4 : Suppression effect on the growth *E. coli* 0157

The suppression effect of lactic acid bacteria from fermented ginseng on the growth of *E. coli* 0157 was investigated. *E. coli* 0157 was inoculated into each 5ml of GAM at the concentration of 10^5 /ml. Each of the culture above was then mixed with each of the lactic acid bacteria listed in Table 9 at the ratio of 1:1 and 1:10 (*E. coli* 0157 to lactic acid bacteria). The mixed culture was then incubated for 20hrs at 37°C and subsequently streaked on a TS agar plate, followed by a incubation at 37°C for 20hrs. The *E. coli* 0157 colonies on the agar plate were counted thereafter.

Table 9

	% inhibition	
	1(harmful bacteria 10^5 /ml) : 1(lactic acid bacteria 10^5 /ml)	1(harmful bacteria 10^5 /ml) : 10(lactic acid bacteria 10^6 /ml)
Control (harmful bacteria only)	0	0
<i>Bifidobacterium</i> KK-1 and harmful bacteria	82	95
<i>Bifidobacterium</i> KK-2 and harmful bacteria	80	85
<i>Bifidobacterium</i> H-1 and harmful bacteria	78	88
<i>Bifidobacterium</i> KK-11 and harmful bacteria	67	85

15

INDUSTRIAL APPLICABILITY

As shown above, fermented products of ginseng containing *Panax quiquefolium* L. and *Panax quinquefolium* L. according to the present invention contain large amount of compound K, Ginsenoside Rh1 and Rh2, and Δ^{20} -ginsenoside Rh2 which are almost not

present, or if any, only in extremely small quantities in unprocessed ginseng. As a result, it can be suggested that the fermented products of ginseng not only have anticancer and anti-allergenic effect but also improve intestinal environment. Furthermore, it has the strong effect of preventing colon cancer, protecting from liver damage and suppressing the growth of harmful intestinal bacteria. Therefore, yogurt containing the fermented product(s) of ginseng as an active component should prove industrially or commercially useful as a yogurt with specific function providing the above mentioned physiological activities of Saponin.




BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Kim Dong Hyun
College of Pharmacy Kyung Hee University
#1 Hocki-Dong, Dongdaemun-Ku
Seoul 130-701
Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Bifidobacterium</i> H-1	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10493
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on May 1, 2003. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: May 6, 2003 

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired; where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.



BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Kim, Dong-Hyun
College of Pharmacy, Kyung-Hee University,
1 Hoeki-Dong, Dongdaemoon-Ku,
Seoul 130-701,
KOREA

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Bifidobacterium</i> KK-1	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10364
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Mar. 22. 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Mar. 29. 2002



¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.



BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Kim, Dong-Hyun

College of Pharmacy, Kyung-Hee University,
1 Hoeki-Dong, Dongdaemoon-Ku,
Seoul 130-701,
KOREA

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : <i>Bifidobacterium</i> KK-2	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10365
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Mar. 22. 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Mar. 29. 2002.



¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.